

Importance and evolution of sexual dimorphism in different families of Collembola (Hexapoda)

José Guadalupe Palacios-Vargas⁽¹⁾ and Gabriela Castaño-Meneses^(1,2)

⁽¹⁾Universidad Nacional Autónoma de Mexico (UNAM), Departamento de Ecología y Recursos Naturales, Facultad de Ciencias, Ecología y Sistemática de Microartrópodos, Ciudad Universitaria 04510, México, D.F. E-mail: jgpv@hp.fciencias.unam.mx ⁽²⁾UNAM, Facultad de Ciencias, Unidad Multidisciplinaria de Docencia e Investigación, Campus Juriquilla, Boulevard Juriquilla 3001, CP 76230, Querétaro, México. E-mail: gcm@hp.fciencias.unam.mx

Abstract – A bibliographic revision of sexual dimorphism in different families of Collembola was performed in order to determine the most common patterns in each group. We list several examples of the sexual dimorphism in different families of Collembola, which illustrate from some of the smallest differences between males and females to major differences between both sexes in some families. We show different cases of sexual dimorphism in *Spinactaletes*, *Sminthurides*, and the remarkable morphology of *Guthriella muskegis*. It seems that most of the cases of strong sexual dimorphism are related to aquatic habitats or behaviour still not fully understood. Species belonging to primarily aquatic families (fresh water or marine water) have the most evolved morphology, where males have clasping antennae to attach the females in order to avoid loosing them with the movement of the water.

Index terms: *Guthriella muskegis*, morphological variations, secondary sexual characters.

Importância e evolução do dimorfismo sexual em diferentes famílias de Collembola (Hexapoda)

Resumo – Uma revisão bibliográfica do dimorfismo sexual em diferentes famílias de Collembola foi feita para determinar os padrões morfológicos mais comuns em cada grupo. Foram listados diversos exemplos de dimorfismo sexual em diferentes famílias de Collembola, que ilustram desde as menores diferenças entre machos e fêmeas até as grandes diferenças entre os sexos em algumas famílias. Foi mostrado o dimorfismo sexual em *Spinactaletes* e *Sminthurides*, e a morfologia marcante de *Guthriella muskegis* foi documentada. É provável que a maioria dos exemplos mais evidentes de dimorfismo sexual estejam relacionados ao habitat aquático ou a padrões de comportamento ainda não compreendidos. As espécies que pertencem às famílias aquáticas primárias (água doce ou água marinha) são as que têm morfologia mais evoluída. Os machos têm antenas preênsis, para agarrar as fêmeas e evitar que se desprendam com o movimento da água.

Termos para indexação: *Guthriella muskegis*, variações morfológicas, caracteres sexuais secundários.

Introduction

Guaranteeing the transference of genetic information to subsequent generations is a fundamental task in all living beings. The success of this process is related to differences in the access to mating, or sexual selection (Darwin, 1871). On mating, competition appears generally between males to be selected by females. The sexual dimorphism in males of Hexapoda appears in different groups, showing complex patterns that include particular behavior for mating.

Nevertheless, even in groups in which there is no copulation, in which the males do not have much contact with females, and the transfer of sperm is done by other mechanisms, there is still some kind of sexual dimorphism.

In Collembola, fecundation is through the transfer of spermatophore, which in some cases is direct, with complex behavior of both sexes (Kozlowski & Aoxiang, 2006), and some species have several modified structures to be able to accomplish that function. Most of these records in Collembola belong to the members of Symphyleona

(Bretfeld, 1971), mainly in *Deuterostomithurus* and in many members of the Sminthurididae (v. gr. *Denisiella*). Nevertheless, there are records of sexual dimorphism in other groups as Isotomidae: *Folsomia* (Ding et al., 2006), *Vertagopus* (Denis, 1928; Goto, 1972; Goloshchapova et al., 2006), *Proisotoma* (Goloshchapova et al., 2006); *Najtia vicaria* (Arlé & Mendonça, 1986); and even in some Entomobryidae: *Seira* (Zeppelini & Bellini, 2006), *Willowsia* (Mari-Mutt, 1981; Zhang et al., 2007), *Orchesella* (Ernsting & Isaaks, 2002) and *Entomobrya* (Ramel et al., 2008); and in the genera *Arrhopalites* (Christiansen & Bellinger, 1996) and *Cyphoderus* (Yoshii, 1990). In *Schoettella* (Hypogastruridae), important differences in the cuticular lipids have been found in each sex (Porco et al., 2004).

In order to determinate the most common patterns in each group, we illustrate the sexual dimorphism found in members of different families in diverse habitats: *Pseudachorutes* (Neanuridae), *Spinactaletes* (Spinactaletidae), *Coenaletes* (Coenaletidae), *Denisiella* (Sminthurididae) and *Guthriella muskegis* (Isotomidae).

Sexual dimorphism determination

Specimens from different habitats were evaluated in order to determine sexual dimorphism: terrestrial (*Pseudachorutes*: Neanuridae; *Guthriella muskegis*: Isotomidae; *Denisiella*: Sminthurididae); primary fresh water (*Sminthurides*: Sminthurididae), littoral (*Spinactaletes*: Spinactaletidae); and associated with hermit crabs (*Actaletes*: Actaletidae). These belonged to the collection of the authors, deposited at the Universidad Nacional Autónoma de México.

Sexual dimorphism in Collembola families

According to the literature reviewed, there are sexual dimorphism records in ten of the 33 families of Collembola. Secondary sexual characters have been recorded in 28 genera. The main examples of sexual dimorphism, found in different families of Collembola, are shown in Table 1.

In the Symphyleona, sexual dimorphism is more evident, and this character affects head and

antennae and the success of spermatophore transfer to the females, regardless of the environment (these modifications occur in edaphic habitats, as well as in aquatic, freshwater or marine littoral). In the case of the terrestrial *Pseudachorutes* (Neanuridae), two ventral organs were observed in the male (Gao & Palacios-Vargas, 2008), and interpreted as sexual dimorphism, which is very clearly symmetrical for both sides of the body. This kind of sexual dimorphism has been observed in other members of the Poduromorpha (as Onychiuridae), and might be common in other families.

In the case of the Isotomidae, the most extraordinary modifications in *Guthriella muskegis* are in the end of the body, in particular the presence of spines and modifications of the setae. Males present ornamentation similar to the patterns found in other groups of Arthropoda, as the Insecta, in order to attract females. These characteristics may only emerge when males are in the reproductive condition and males very similar to females can be found, even in the adult phase, without modifications. In modified males, the setae look like leaves and they develop abdominal spines, which are not present in the females; even the antennae are modified (Figure 1). There are also records of modifications of the antennae in other species of Isotomidae, their function are still unknown. The specimens of *G. muskegis* are from Cerro Fábrega, Bocas del Toro, Panama.

For the Entomobryidae, the main sexual differences are in the color patterns, and in the modification of mucro and genital aperture.

In the case of the species members of the littoral, *Spinactaletes* spp. (Spinactaletidae), the modifications are the presence of a spur in the tibiotarsus III and also clasping antennae in the males.

For the peculiar case of *Coenaletes caribaeus* Bellinger (Coenaletidae), which lives between the shell and the hermit crab, we studied the differences of males and females (Palacios-Vargas et al., 2000). The modifications include the presence of clasping antennae with spines and also some spines on thorax and abdomen, although their function are still unknown.

Table 1. Sexual dimorphism among Collembola.

| Taxa | Characteristics | References |
|---|--|---|
| Neanuridae <i>Pronura pomorskii</i> Smolis & Deharveng | Fore legs structure | Smolis & Deharveng (2006) |
| Hypogastruridae <i>Hypogastrura assimilis</i> (Krasbauer) <i>Schoettella ununguiculata</i> (Tullberg) | Body length Cuticular lipids | Folker-Hansen et al. (1996) Porco et al. (2004) |
| Isotomidae <i>Agrenia</i> spp. <i>Anurophorus serratus</i> Deharveng <i>Archisotoma besselsi</i> (Pack.) <i>Axelsonia tubifera</i> Strenzke <i>Dimorphotoma porcella</i> Ellis <i>Folsomia fimetaria</i> (L.) <i>Jestella siva</i> Najt <i>Najtia vicaria</i> (Arlé) <i>Rhodanella minos</i> (Denis) <i>Vertagopus pseudocinereus</i> Fjellberg | Male antenna elongated with more developed 2 nd and 3 rd segments. Curved sensilla occur on the ventromedial side of the three last antennal segments of males. Male antenna elongated with more developed 2 nd and 3 rd segments. Abdominal shortened macrosetae forming a "brush". Male antenna elongated and broadened with more developed 2 nd and 3 rd segments. Male antenna elongated and broadened with more developed 2 nd and 3 rd segments. Male abdominal setae longer and thicker than in females. Body length Male antenna elongated and broadened with more developed 2 nd and 3 rd segments. Male abdominal setae longer and thicker than in females. Male antenna elongated, Ant. IV horn-like curved, lateral cephalic macrosetae, Abd. IV with crowbar-like structures. Presence of a pair of very long, out-standing, curved horn-like setae, one on either side of the head, and by the presence of stouter antennae and modified setae on some of the abdominal segments. Males 2 nd antennal segment curved and with slightly serrated setae. Shortened macrosetae forming a "brush". | Strenzke (1958) Deharveng (1976) Denis (1928) Fjellberg (1986) Ellis (1976) Folker-Hansen et al. (1996) Najt (1977) Arlé & Mendonça (1986) Goto (1972) Goloshchapova et al. (2006) |
| Actaletidae <i>Spinactaletes</i> spp. | Length of male antenna, form of Ant. IV, chaetotaxie of Ant. II, III and IV, wide of Ant. I. Prefemoral and protibiotarsal chaetotaxie of males. | Soto-Adames (1988) |
| Entomobryidae <i>Entomobrya atrocincta</i> Schött <i>Seira mantis</i> Zeppelini & Bellini <i>S. raptora</i> Zeppelini & Bellini <i>Orchesella cincta</i> (L.) <i>Willowsia jacobsoni</i> (Börner) | Body color pattern Modifications in the fore legs Modifications in the fore legs Body color pattern Body color pattern and body length | Ramel et al. (2008) Zeppelini & Bellini (2006) Zeppelini & Bellini (2006) Mari-Mutt (1981) Mari-Mutt (1981) |
| Paronellidae <i>Cyphoderus albinus</i> Nicolet | Females with bidentate muco | Yoshii (1990) |
| Coenaletidae <i>Coenaletes caribaeus</i> Bellinger | Fourth antennal segment in male with 2 subapical organs in separate minute pegs and 8 th thin, uniform blunt setae in apical half. | Palacios-Vargas et al. (2000) |
| Sminthurididae <i>Denisiella diomedesi</i> Palacios-Vargas <i>Debouttevillea marina</i> Murphy <i>Jeannenotia stachi</i> Jeannenot <i>Sminthuridia sphaeridioides</i> (Murphy) <i>Yosiides himachal</i> (Yosii) | Antenna of males modified as clasping organ Second and third antennal segment highly modified as clasping organ in males. Males second antennal segment modified in a clasping organ. Males second antennal segment modified in a clasping organ. Males second antennal segment modified in a clasping organ. | Palacios-Vargas (2007) Massoud & Betsch (1972), Eberhard (2004) Massoud & Betsch (1972) Massoud & Betsch (1972), Eberhard (2004) Massoud & Betsch (1972), Eberhard (2004) |
| Arrhopalitidae <i>Arrhopalites jay</i> Christiansen & Bellinger <i>Arrhopalites lacuna</i> Christiansen & Bellinger | Presence of a tunica and an unguicular filament on the posterior foot of female Number of antennal subsegments and the fact that the male seta e3 differs strikingly from the female | Christiansen & Bellinger (1996) Christiansen & Bellinger (1996) |
| Bourletiellidae <i>Bovicornia greensladei</i> Massoud & Delamare Deboutteville <i>Deuterostminthurus</i> spp. | Males second antennal segment modified as clasping organ Body and head length, longer antennae in males. Female with subanal appendages | Eberhard (2004) Palacios-Vargas & González (1995) |




Figure 1. Female and male of *Guthriella muskegis*. Dorsal view of female (A), dorsal view of male (B), last abdominal segments in female (C), and last abdominal segments in male (D).

Conclusions

1. The habitat can be a very important factor determining the presence of several sexual characters.
2. Fresh water and marine littoral springtails present more important morphological modifications to facilitate sperm transfer from the male to the female.
3. In terrestrial species, the most extraordinary case is that of *G. muskegis* which ornamentation is similar to the patterns found in other groups of Arthropoda.

Acknowledgements

To Dr. Diomedes Quintero (Universidad de Panamá), for the donation of material of *Guthriella muskegis*; to Dr. Silvia Espinosa-Matías (Fac. Ciencias, UNAM), for scanning microphotographs; to Dr. Douglas Zeppelini

(Universidad Federal da Paraíba, Brazil) and Mirna Samaniego (Smithsonian Tropical Research Institute, Panama), for review of the Portuguese abstract.

References

- ARLÉ, R.; MENDONÇA, M.C. de. Descrição do macho de *Isotoma vicaria* Arlé, 1959 com proposta de um novo gênero (Collembola, Isotomidae). *Revista Brasileira de Entomologia*, v.30, p.1-4, 1986.
- BRETFELD, G. Das Paarungsverhalten europäischer Bourletiellini (Sminthuridae). *Revue d'Ecologie et de Biologie du Sol*, v.8, p.145-153, 1971.
- CHRISTIANSEN, K.; BELLINGER, P. Cave *Arrhopalites*: new to science. *Journal of Cave and Karst Studies*, v.58, p.168-180, 1996.
- DARWIN, C. **The descent of man and selection in relation to sex**. London: John Murray, 1871. 424p.

- DEHARVENG, L. Présence d'un caractère sexuel secondaire chez *Anuophorus serratus* n. sp. (Collembola: Isotomidae). **Nouvelle Revue de Entomologie**, v.6, p.109-111, 1976.
- DENIS, J.R. Sur deux collemboles de la Somalie Italienne. Le dimorphisme sexuel de *Vertagopus minos* n. sp. **Bollettino della Società Entomologica Italiana**, v.60, p.1-6, 1928.
- DING, Y.F.; HUANG, C.H.; CHEN, J.X. A new species group in the genus *Folsomia* (Collembola: Isotomidae), with a new species from the Three Gorges Region, China. **Entomological News**, v.117, p.553-558, 2006.
- EBERHARD, W.G. Rapid divergent evolution of sexual morphology: comparative tests of antagonistic coevolution and traditional female choice. **Evolution**, v.58, p.1947-1970, 2004.
- ELLIS, W.N. Autumn fauna of Collembola from Central Crete. **Tijdschrift voor Entomologie**, v.119, p.221-326, 1976.
- ERNSTING, G.; ISAAKS, J.A. Gamete production and sexual size dimorphism in an insect (*Orchesella cincta*) with indeterminate growth. **Ecological Entomology**, v.27, p.145-151, 2002.
- FJELLBERG, A. Revision of the genus *Agrenia* Börner, 1906 (Collembola: Isotomidae). **Entomologica Scandinavica**, v.17, p.93-106, 1986.
- FOLKER-HANSEN, P.; KROGH, P.H.; HOLMSTRUP, M. Effect of dimethoate on body growth of representatives of the soil-living mesofauna. **Ecotoxicology and Environmental Safety**, v.33, p.207-216, 1996.
- GAO, Y.; PALACIOS-VARGAS, J.G. A new species of genus *Pseudachorutes* Tullberg, 1871 (Collembola: Neanuridae: Pseudachorut) from China. **Zootaxa**, v.1895, p.53-58, 2008.
- GOLOSHCHAPOVA, N.P.; POTAPOV, M.B.; CHERNOVA, N.M. Sexual behaviour in Isotomidae (Collembola). **Pedobiologia**, v.50, p.111-116, 2006.
- GOTO, H.E. Some observations on the biology and taxonomy of *Proisotoma stachi* Goto, 1957 (Insecta, Collembola: Isotomidae). **Journal of Natural History**, v.6, p.195-202, 1972.
- KOZLOWSKI, M.W.; AOXIANG, S. Ritual behaviors associated with spermatophore transfer in *Deuterostimnthus bicinctus* (Collembola: Bourletiellidae). **Journal of Ethology**, v.24, p.103-109, 2006.
- MARI-MUTT, J.A. Redescription of *Willowsia jacobsoni* (Börner), an entomobryid with conspicuous sexual dimorphism (Insecta: Collembola). **Journal of Agriculture of the University of Puerto Rico**, v.65, p.361-373, 1981.
- MASSOUD, Z.; BETSCH, JM. Étude sur les insectes collemboles. II. Les caractères sexuels secondaires des antennes des Symphypléones. **Revue d'Ecologie et de Biologie du Sol**, v.9, p.55-97, 1972.
- NAJT, J. Un nouveau genre de Collembole Isotomidae du Nepal: *Jestella siva* n.g. n.sp., **Nouvelle Revue de Entomologie**, v.7, p.363-375, 1977.
- PALACIOS-VARGAS, J.G. A new species of *Denisiella* (Collembola: Sminthurididae) from Panama and new records for *D. sexpinnata* (Denis, 1938). **Zootaxa**, v.1637, p.63-68, 2007.
- PALACIOS-VARGAS, J.G.; CUTZ, L.Q.; MALDONADO, C. Redescription of the male of *Coenaletes caribaeus* (Collembola: Coenaletididae) associated with hermit crab (Decapoda: Coenobitidae). **Annals of the Entomological Society of America**, v.93, p.194-197, 2000.
- PALACIOS-VARGAS, J.G.; GONZÁLEZ, V. Two new species of *Deuterostimnthus* (Bourletiellidae), epiphytic Collembola from the Neotropical region with a key for the American species. **Florida Entomologist**, v.78, p.286-294, 1995.
- PORCO, D.; DEHARVENG, L.; GERS, C.H. Sexual discrimination with cuticular lipids in *Schoettella ununguiculata* (Tullberg, 1869) (Collembola: Hypogastruridae). **Pedobiologia**, v.48, p.581-583, 2004.
- RAMEL, G.; BAQUERO, E.; JORDANA, R. Biodiversity of the Collembola fauna of wetland Kerkini (N. Greece), with description of the sexual dimorphism of *Entomobrya atrocincta* Schött 1896 (Collembola: Entomobryomorpha). **Annales de la Société Entomologique de France**, v.44, p.113-128, 2008.
- SHINE, R. Ecological causes for the evolution of sexual dimorphism: a review of the evidence. **The Quarterly Review of Biology**, v.64, p.419-461, 1989.
- SMOLIS, A.; DEHARVENG, L. A new species of *Pronura* Delamare Deboutteville, 1953 from North of Vietnam (Collembola: Neanuridae: Neanurinae). **Annales Zoologici**, v.56, p.443-448, 2006.
- SOTO-ADAMES, F.N. Revisión de la familia Actaletidae Börner, 1902 (Insects: Collembola). **Caribbean Journal of Science**, v.24, p.161-196, 1988.
- STRENZKE, K. *Axelsonia tubifera* n.sp., ein neuer arthropleoner Collembole mit Geschlechtsdimorphismus aus der brasilianischen Mangrove. **Acta Zoologica Cracoviensia**, v.2, p.607-618, 1958.
- YOSHII, R. Miscellaneous notes on the Collembola of Macaronesia. **Contributions from the Biological Laboratory of Kyoto University**, v.27, p.535-540, 1990.
- ZEPPELINI, D.; BELLINI, B. Two *Seira* Lubbock 1869 (Collembola, Arthropleona, Entomobryidae) new to science, with remarkable secondary sexual characters. **Zootaxa**, v.1185, p.21-35, 2006.
- ZHANG, F.; PALACIOS-VARGAS, J.G.; CHEN, J.-X. The genus *Willowsia* and its Mexican species (Collembola: Entomobryidae). **Annals of the Entomological Society of America**, v.100, p.36-40, 2007.

Received on September 9, 2008 and accepted on July 3, 2009